

**“A COMPARATIVE STUDY OF  
DEXMEDETOMIDINE AND ESMOLOL IN  
INDUCED HYPOTENSION FOR FUNCTIONAL  
ENDOSCOPIC SINUS SURGERY”**

**Dissertation submitted in partial fulfillment of**

**M.D. DEGREE EXAMINATION**

**M.D. ANAESTHESIOLOGY- BRANCH X**

**CHENGALPATTU MEDICAL COLLEGE, CHENGALPATTU**



**THE TAMILNADU DR. M.G.R. MEDICAL UNIVERSITY**

**CHENNAI, TAMILNADU.**

**APRIL 2015**

## **CERTIFICATE**

This is to certify that this dissertation titled “**comparative study of dexmedetomidine and esmolol in induced hypotension for functional endoscopic sinus surgery**” has been prepared by Dr .B. Rajesh under my supervision in the department of anesthesiology, chengalpattu medical college and hospital, chengalpattu during the academic period 2012-2015 and is being submitted to the Tamil Nadu Dr. M.G.R. Medical university, Chennai in partial fulfillment of the university regulation for the award of the Degree of Doctor of Medicine (Branch –X MD anesthesiology) and his dissertation is a bonafide work.

**Prof Dr.RAJA BILLI GRAHAM,MS.,**  
DEAN  
Chengalpattu medical college&  
Hospital  
Chengalpattu.

**Prof.Dr. SUGANTHARAJ**  
**ANURADHA M.D.D.A**  
Professor & HOD,  
Department of Anaesthesiology  
Chengalpattu Medical College  
Chengalpattu.

## **DECLARATION**

I, **Dr.RAJESH .B**, solemnly declare that the dissertation “**A COMPARATIVE STUDY OF DEXMEDETOMIDINE AND ESMOLOL IN INDUCED HYPOTENSION FOR FUNCTIONAL ENDOSCOPIC SINUS SURGERY**” is a bonafide work done by me in the Department of Anaesthesiology, Chengalpattu Medical College & Hospital, Chengalpattu, after getting approval from the Ethical committee under the able guidance of **Prof.Dr. SUGANTHARAJ ANURADHA M.D.,D.A.**, Professor & HOD, Department of Anaesthesiology, Chengalpattu Medical College, Chengalpattu.

Place : Chengalpattu

Date :

**(Dr.RAJESH.B)**

## **ACKNOWLEDGEMENT**

I am extremely thankful to **Dr.RAJA BILLI GRAHAM, M.S.,** Dean, Chengalpattu Medical College, for her permission to carry out this study.

I am immensely grateful to **PROF. Dr.SUGANTHARAJ ANURADHA, M.D.,D.A.,** Professor and Head of the Department, Department of Anaesthesiology, for his concern and support in conducting this study.

I am very grateful to express my sincere gratitude to the Additional Professors **Dr. VALLI SATHYAMOORTHY, M.D.D.A., Dr.J.REVATHY M.D.D.A.,** Department of Anaesthesiology, for their constant motivation and valuable suggestions.

I am extremely grateful to the Assistant Professor **Dr.M.RAVIKUMAR M.D., D.A,** for his guidance and expert advice in carrying out this study.

I am thankful to the Institutional Ethical Committee for their guidance and approval for this study.

I would like to thank the Assistant Professors and residents of the Department of ear ,nose and throat for their unwavering support for this study.

I am thankful to all my colleagues and friends for their help and advice in carrying out this dissertation.

Last but not least, I thank all the patients for willingly submitting themselves for this study.

**INSTITUTIONAL ETHICS COMMITTEE**  
**CHENGALPATTU MEDICAL COLLEGE , CHENGALPATTU**  
**APPROVAL OF ETHICAL COMMITTEE**

To

Dr.B.Rajesh  
Post Graduate  
Dept of Anaesthesia

Dear Dr.

The Institutional Ethical Committee of Chengalpattu Medical College reviewed and discussed your application to conduct the clinical / dissertation work entitled

INDUCED HYPOTENSION FOR FESS (FUNCTIONAL ENDOSCOPIC SINUS SURGERY) -  
A COMPARATIVE STUDY OF DEXMEDITOMEDINE VS ESMOLOL

On 13.11.2013

The following documents reviewed

- a. Trial protocol, dated \_\_\_\_\_ version no
- b. Patient information sheet and informed consent form in English and / or vernacular language.
- c. Investigators Brochure, dated \_\_\_\_\_ version
- d. Principal Investigators current CV
- e. Investigators undertaking

The following members of the Ethics committee were present at the meeting held on

Date 13.11.2013 Time 12.00 Noon Place Chengalpattu Medical College

Approved  Chairman Ethics Committee

 13/11/13 Member secretary of Ethics Committee.

Name of each member with designation

Clinical Members

1. Dr.G.Raja Billy Graham MS.,  
Prof & HOD of Surgery, CHMC
2. Dr.K.Srinivasagalu MD.,  
Prof & HOD of Medicine, CHMC

*G.R. Graham*  
*[Signature]*

Biological Scientist

3. Dr.K.Baskaran MD.,  
Asso Prof of Pharmacology, CHMC

Non Clinical Members

4. Dr.P.Parasakthi MD  
Prof & HOD of Forensic Medicine,CHMC
5. Member from Nongovernmental  
Voluntary Organisation : Mr.P.Durairaj
6. Philosopher : Mr.K.S.Ramprasad
7. Lawyer : Lr. I. M. Karimala Basha
8. Layperson : Mr.Dilli

*[Signature]*

*W. Dilli*

We approve the clinical trial to be conducted in its presented form

The Institutional Ethics Committee expects to be informed about the progress of the study and any SAE occurring in the course of the study, any changes in protocol and patient information / informed consent and asks to provide copy of final report.

Yours sincerely

*[Signature]*  
13/11/13  
Member secretary, Ethics Committee

Turnitin Document Viewer - Mozilla Firefox

https://www.turnitin.com/dv?s=1&o=457015623&u=1030927302&student\_user=1&lang=en\_us&

The Tamil Nadu Dr.M.C.R.Medical ...TNMGRMU EXAMINATIONS - DJE 15-A...

Originality

GradeMark

PeerMark

Dissertation

BY 201220202.MD

turnitin

13%SIMILAR

OUT OF 0

Match Overview

1

Xin, H., and W. Jlanan. ...

Publication

1%

2

Hanna Phan. "Clinical ...

Publication

1%

3

Yang, C.-Y., C.-F. Cre ...

Publication

1%

4

Sher I Kashmir, Imtiaz ...

Publication

1%

5

Castro, Diogo S., Pedr...

Publication

1%

INTRODUCTION

6

Functional endoscopic sinus surgery is one of the commonly performed

surgeries

Induced hypotension is a method employed in functional endoscopic

sinus surgery to reduce the blood loss and to improve visibility of the surgical

field induced hypotension also helps in reducing blood transfusion during surgery

Induced hypotension is also known as controlled hypotension

Go Back

Download

Print

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770

771

772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

846

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875

876

877

878

879

880

881

882

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926

927

928

929

930

931

932

933

934

935

936

937

938

939

940

941

942

943

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

959

960

961

962

963

964

965

966

967

968

969

970

971

972

973

974

975

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

992

993

994

995

996

997

998

999

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268

1269

1270

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

1303

1304

1305

1306

1307

1308

1309

1310

1311

1312

1313

1314

1315

1316

1317

1318

1319

1320

1321

1322

1323

1324

1325

1326

1327

1328

1329

1330

1331

1332

1333

1334

1335

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

1348

1349

1350

1351

1352

1353

1354

1355

1356

1357

1358

1359

1360

1361

1362

1363

1364

1365

1366

1367

1368

1369

1370

1371

1372

1373

1374

1375

1376

1377

1378

1379

1380

1381

1382

1383

1384

1385

1386

1387

1388

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

1400

1401

1402

1403

1404

1405

1406

1407

1408

1409

1410

1411

1412

1413

1414

1415

1416

1417

1418

1419

1420

1421

1422

1423

1424

1425

1426

1427

1428

1429

1430

1431

1432

1433

1434

1435

1436

1437

1438

1439

1



## **CONTENTS**

<b>S.NO</b>	<b>TOPIC</b>	<b>PAGE NO</b>
<b>1</b>	<b>INTRODUCTION</b>	<b>1</b>
<b>2</b>	<b>AIM</b>	<b>2</b>
<b>3</b>	<b>HYPOTENSIVE ANESTHESIA</b>	<b>3</b>
<b>4</b>	<b>FUNCTIONAL ENDOSCOPIC SINUS SURGERY</b>	<b>18</b>
<b>5</b>	<b>PHARMACOLOGY OF ESMOLOL</b>	<b>22</b>
<b>6</b>	<b>PHARMACOLOGOLGY OF DEXMEDETOMIDINE</b>	<b>30</b>
<b>7</b>	<b>REVIEW OF LITERATURE</b>	<b>37</b>
<b>8</b>	<b>MATERIALS AND METHODS</b>	<b>48</b>
<b>9</b>	<b>RESULTS AND OBSERVATION</b>	<b>57</b>
<b>10</b>	<b>DISCUSSION</b>	<b>91</b>
<b>11</b>	<b>SUMMARY</b>	<b>98</b>
<b>12</b>	<b>CONCLUSION</b>	<b>100</b>
<b>13</b>	<b>BIBLIOGRAPHY</b>	
<b>14</b>	<b>PROFORMA</b>	
<b>15</b>	<b>MASTER CHART</b>	

## **ABBREVIATION**

Paco <sub>2</sub>	-	partial pressure of carbon di oxide in arterial blood
Fio <sub>2</sub>	-	inspired oxygen concentration
ECG	-	electrocardiogram
ETCO <sub>2</sub>	-	end tidal carbon di oxide
CSF	-	cerebro spinal fluid
COPD	-	chronic obstructive pulmonary disease
MAP	-	mean arterial pressure
FESS	-	functional endoscopic sinus surgery
ASA	-	American society of anaesthesiologist

# ABSTRACT

## OBJECTIVE

To compare the effectiveness and advantages of using dexmedetomidine and esmolol in induced hypotension for functional endoscopic sinus surgery.

## METHODS

Fifty patients belonging to ASA 1 planned for functional endoscopic sinus surgery were randomly divided into two groups. Dexmedetomidine group received a bolus dose of 1 microgram/kg over 10 minutes followed by infusion at a rate of 0.4-0.8 microgram /kg/hr. Esmolol group received a bolus dose of 1 mg/kg over one minute followed by infusion at a rate of 0.4-0.8mg/kg/hr. The target mean arterial pressure was 55-65 mm hg. General anaesthesia was maintained with 60% N<sub>2</sub>O:O<sub>2</sub>. Induction dose of propofol, haemodynamic variables (blood pressure and heart rate), intra operative serum cortisol, quality of surgical field, emergence time, post anaesthesia recovery score(modified Aldrete score >9), sedation score at 15, 30 and 60 minutes, time for rescue analgesia were recorded.

## RESULT

Dose of propofol was lower in dexmedetomidine group than esmolol group. Heart rate and blood pressure were comparable in both groups intra operatively, when target mean arterial was achieved. Quality of surgical field and

intra operative serum cortisol level's were comparable in both group's. Emergence time and time to achieve aldrete score  $> 9$  were higher in dexmedetomidine group than esmolol group. Sedation score's were higher in dexmedetomidine group than esmolol group at 15 ,30 minute's post operatively. Time for rescue analgesia was prolonged in dexmedetomidine group than esmolol group.

## CONCLUSION

The use of dexmedetomidine in induced hypotension for functional endoscopic sinus surgery has additional advantage of sedative ,analgesic and reduced requirement of anaesthetic agent's than esmolol.

## **INTRODUCTION**

Functional endoscopic sinus surgery is one of the commonly performed Surgeries.

Induced hypotension is a method employed in functional endoscopic sinus surgery to reduce the blood loss and to improve visibility of the surgical field. Induced hypotension also helps in reducing blood transfusion during surgery.

Induced hypotension is also known as controlled hypotension (or) deliberate hypotension (or) hypotensive anesthesia. Induced hypotension also prevents progressive stretching of the vessel wall during vascular surgery, which enables easy suturing and clipping in aortic surgery , AV malformations and aneurysmal surgeries.

Esmolol is an ultrashort acting beta blocker and dexmedetomidine is a highly selective alpha 2 adrenergic agonist .

Hence,the above study was done in the department of anaesthesiology, chengalpattu medical college,Chengalpattu.

## **AIM**

The aim of the study was to compare the effectiveness and advantages of using dexmedetomidine and esmolol in induced hypotension for functional endoscopic sinus surgery.

## **HYPOTENSIVE ANESTHESIA**

Induced hypotension is a method employed to improve the quality of the surgical field and to minimize the complications associated with blood transfusion.

### **HISTORY**

- Induced hypotension was first proposed by Harving cushing for neurosurgery in 1917.
- Gardner was the first to practice induced hypotension in surgery by arteriotomy.
- Griffith and gillies used high spinal anesthesia for producing Hypotension.
- Enderby used ganglionic blockers for induced hypotension and he also insisted on the importance of posture in induced hypotension.
- Sodium nitroprusside was first used hypotensive anesthesia by Moraca in 1962.
- Dimant used cardiac pacemakers for hypotensive anesthesia in 1967.
- Nitroglycerine and labetalol were used for induced hypotension in 1970.

## **PHYSIOLOGY OF INDUCED HYPOTENSION**

During hypotensive anesthesia adequate blood flow should be maintained to vital organs in order to prevent end organ damage. Blood flow to many vital organs such as heart, brain, kidneys and liver is autoregulated, which enables them to be perfused over a wide range of perfusion pressures.

### **OBJECTIVES**

- Maintain systolic blood pressure between 80-90mm hg.
- Maintain mean arterial pressure between 50-65mm hg.
- Maintain mean arterial pressure 30% of baseline value in hypertensive patients.
- A close communication between surgeon and anesthetist is necessary in order to maintain optimal surgical conditions without compromising end organ perfusion.
- Critical perfusion pressure varies for different organs and individuals.

### **CENTRAL NERVOUS SYSTEM**

- Cerebral blood flow is autoregulated between 50-150mm hg.



- During hypotensive anesthesia when cerebral blood flow reduces more the cerebral metabolic rate, cerebral ischemia develops.
- Volatile anesthetics used in hypotensive anesthesia attenuate the cerebral autoregulation.
- Vasodilators used in hypotensive anesthesia attenuate the cerebral autoregulation.
- Head end elevation in hypotensive anesthesia also reduces the cerebral blood flow.

## **RESPIRATORY SYSTEM**

- Ventilation perfusion mismatch can occur during hypotensive anesthesia due to use of head end elevation.
- Increase in  $paco_2$  can be due to inhibition of hypoxic pulmonary vasoconstriction leading to increase in intra pulmonary shunting.
- In order to avoid hypoxia higher  $fio_2(0.4-0.5)$  should be used during hypotensive anesthesia.

## **CARDIOVASCULAR SYSTEM**

- Coronary blood flow decreases during hypotensive anesthesia, but corresponding decrease in myocardial oxygen demand prevents development of myocardial ischemia in healthy individuals.

- Hypotensive anesthesia, when used in patients with coronary artery occlusion can cause redistribution of blood flow leading to myocardial infarction.

### **RENAL SYSTEM**

- Hypotensive anesthesia causes decrease in glomerular filtration rate leading to oliguria in intraoperative period.
- Adequate hydration should be maintained in order to avoid postoperative renal dysfunction in hypotensive anesthesia.

### **HEPATIC SYSTEM**

- Hepatic functions are not affected by hypotensive anesthesia.

## **INDICATIONS**

- 1) Neurosurgery-AV malformations, cerebral aneurysm surgery, vascular tumour surgery, trans sphenoidal hypophysectomy.
- 2) Major orthopaedic surgery-total hip and knee replacement, spine surgery.
- 3) ENT surgery-functional endoscopic sinus surgery, middle ear surgery.
- 4) Head and neck surgery-faciomaxillary tumors, radical neck dissections, larygectomy.
- 5) Ophthalmic surgery-orbital surgery , vitrectomy, intraocular tumors
- 6) Major pelvic and gynaecological surgeries
- 7) Plastic and reconstructive surgery
- 8) Jehovahs witness

## **CONTRAINDICATIONS**

- 1) Ischemic heart disease
- 2) Cerebrovascular disease
- 3) Hepatic dysfunction
- 4) Renal dysfunction
- 5) Severe respiratory disease
- 6) Diabetes mellitus
- 7) Uncontrolled hypertension
- 8) Bleeding diathesis
- 9) Allergy to hypotensive agents
- 10) Severe anemia
- 11) Lack of knowledge of the technique
- 12) Lack of adequate monitoring
- 13) Patient refusal

## **METHODS OF HYPOTENSIVE ANESTHESIA**

### **PHYSICAL MEASURES**

- **Head end elevation**-Elevation of head end of the patient helps to prevent blood loss during surgery, but head end elevation is associated with the risk of air embolism.
- Head end elevation causes pooling of blood in the lower extremities leading to decrease in venous return to the heart and cardiac output.
- **Positive pressure ventilation**-Positive pressure ventilation reduces venous return to the heart, which can be further reduced by prolonging the inspiratory time and by applying positive end expiratory pressure.

### **PHARMACOLOGICAL AGENTS**

#### **Characteristics of a ideal hypotensive agent**

- Easy to administer
- Rapid onset of action
- Hypotensive effect should disappear rapidly on discontinuation
- Rapid elimination
- No toxic metabolites

- Negligible effect on vital organs
- Predictable effect
- Dose dependant effect

## **GANGLIONIC BLOCKING AGENTS**

- They produce hypotension by blocking both sympathetic and parasympathetic outflow from autonomic ganglia.
- **Pentamethonium and hexamethonium** were initially used in hypotensive anesthesia.
- **Pentolinium** is a highly potent agent, devoid of tachycardia due to effective sympathetic blocking action.
- **Trimethaphan** has additional direct vasodilating property and histamine releasing property , which enhances it's hypotensive effect.
- **Trimethaphan** has very short duration of action of 1-2 min's and it is metabolised by plasma cholinesterase.

## **ALPHA ADRENERGIC BLOCKERS**

- Alpha blockers produce hypotension by blocking alpha 1 receptors in blood vessels leading to vasodilatation.

- **Phentolamine** is a short acting alpha blocker used in hypotensive anesthesia.
- **Phentolamine** has rapid onset of action ,it's plasma level reaches a peak within 2 min's.
- **Phentolamine** is usually administered in bolus doses of 2 -5 mg.
- **Labetalol** is a combined alpha and beta blocker.
- **Labetalol** has more potent action on beta receptors when compared with alpha receptors.
- **Labetalol** has rapid onset of action with peak plasma concentrations reaching within 5 min.
- Advantages of labetalol include lack of rebound hypertension, lack of tachycardia, tachyphylaxis.
- Contraindications for labetalol are patients with heart block, bradycardia, conduction block, congestive cardiac failure and hypovolemia.

## **VASODILATORS**

### **1) SODIUM NITROPRUSSIDE**

- Mechanism of action-sodium nitroprusside acts by production of nitric oxide.Nitric oxide diffuses across the cell membrane and

stimulates guanylate cyclase, which inhibits intracellular release of calcium leading to relaxation.

- Nitroprusside has rapid onset of action within 1 min.
- Dose of nitroprusside is 0.5-10 microgram/kg/min.
- Metabolism-Cyanide released from nitroprusside combines with thiosulfate to form thiocyanate in the liver in the presence of enzyme rhodanase. Cyanide also combines with hydroxycobalamine to form cyanocobalamine.
- Complication-Important complication associated with nitroprusside is cyanide toxicity.
- Cyanide can combine with cytochrome oxidase in the electron transport chain leading to histotoxic hypoxia.
- **Manifestations of cyanide toxicity**
  - 1) Resistance to high infusion doses of nitroprusside
  - 2) Metabolic acidosis
  - 3) Elevated lactate levels
  - 4) Hypotension not responding to fluids and vasopressors
  - 5) Cardiovascular collapse



- **Management of cyanide toxicity**

- 1) Sodiumthiosulphate-it converts cyanide to thiocyanate and enables its excretion. It is given as a bolus dose of 30mg/kg followed by infusion at a rate of 60 mg/kg/hr.
- 2) Hydroxocobalamine-it prevents increase in concentration of cyanide in erythrocytes. It is given as a bolus dose of 50mg/kg followed by infusion at a rate of 100 mg/kg/hr.
- 3) Sodium nitrite (or) amyl nitrite-it produces methemoglobin, which combines with cyanide to form cyanomethemoglobin. It is administered at a dose of 5mg/kg intravenously.

## **NITROGLYCERINE**

- Nitroglycerine is predominantly a venodilator, it reduces venous return to the heart and produces hypotension.
- Dose of nitroglycerine is 0.5-10 microgram/kg/min.
- It has rapid onset of action and a short half life of about 2 min.
- Advantages of nitroglycerine is that it does not produce any toxic metabolites and it does not produce rebound hypertension.

## **HYDRALAZINE**

Hydralazine is predominantly a arterial vasodilator used to produce hypotension.

## **NICARDIPINE**

- Nicardipine is a calcium channel blocker used in induced hypotension.
- Nicardipine is given intravenously and it has a distribution half life of 14 min.
- Nicardipine should be cautiously used in patients with renal dysfunction .

## **ADENOSINE**

- Adenosine is a potent vasodilator used in induced hypotension.
- Adenosine has a very short half life of 10-20 sec.
- Adenosine causes decrease in peripheral vascular resistance without causing increase in heart rate.
- Adenosine is infused at a rate of 60-300 microgram/kg/min.
- Advantages of adenosine is that it does not produce any rebound hypertension.

- Disadvantages –It can block the conducting system of heart leading to bradycardia. It can cause coronary steal phenomenon. It can cause accumulation of uric acid in patient with gout.

### **PROSTAGLANDIN E1**

- Prostaglandin E1 has positive inotropic action on the heart and it causes a decrease in systemic vascular resistance.
- Prostaglandin E1 has antiarrhythmic action.
- Prostaglandin E1 helps in maintaining cerebral autoregulation.
- Prostaglandin E1 used in induced hypotension helps in increasing renal blood flow and urine formation.

### **INHALATIONAL AGENTS**

- Inhalational agents like halothane, isoflurane, sevoflurane, desflurane can be used in induced hypotension.
- Halothane is a direct myocardial depressant and it sensitizes the heart to arrhythmogenic action of adrenaline.
- Isoflurane produces hypotension by decreasing the systemic vascular Resistance.
- Sevoflurane and desflurane are newer inhalational agents which produce hypotension without significant myocardial depression.

## **BETA BLOCKERS**

- Beta blockers produce hypotension by decreasing the myocardial contractility and cardiac output.
- Propranolol is a non selective beta blocker used in hypotensive anaesthesia. It is administered at a dose of 0.5mg intravenously. Propranolol can produce bronchospasm in patients with bronchial asthma.
- Metoprolol is a selective beta1 blocker used in induced hypotension.
- Esmolol is a ultrashort acting selective beta 1 blocker used in induced hypotension.

## **REMIFENTANIL**

- Remifentanyl is a ultra short acting opioid used in hypotensive anesthesia.
- Remifentanyl has a ester group and hence it is metabolized by plasma and RBC esterases.
- Remifentanyl has a short context sensitive half life of about 4 minutes.
- Remifentanyl is given at a loading dose of 1microgram/kg/min.
- Remifentanyl can also be used to blunt the sympathetic response associated with laryngoscopy and intubation.

## **MONITORING IN HYPOTENSIVE ANESTHESIA**

- Invasive blood pressure-Intra arterial blood pressure monitoring is gold Standard during hypotensive anesthesia for accurate measurement of blood pressure.
- ECG-Ecg is mandatory during hypotensive anesthesia to detect any compromise in coronary blood flow.
- Pulse oxymeter-pulse oxymeter is used to make changes in FIO<sub>2</sub> due to ventilation perfusion mismatch occurring during hypotensive anesthesia.
- ETCO<sub>2</sub>-Etco<sub>2</sub> is essential as it can be useful in estimation of blood PaCO<sub>2</sub> during hypotensive anesthesia.
- Temperature-Temperature should be monitored to detect hypothermia during hypotensive anesthesia.
- Urine output-urine output should be monitored to assess the adequacy of renal perfusion during hypotensive anesthesia.

## **FUNCTIONAL ENDOSCOPIC SINUS SURGERY**

- Functional endoscopic sinus surgery is a technique to gain access to paranasal sinuses by minimal invasive methods .
- The main aims of sinus surgery is to provide drainage to the sinus cavities and to obtain tissues from sinus cavity for pathological examination.

### **INDICATIONS**

- 1) Chronic sinusitis-The most common indication of endoscopic sinus is chronic sinusitis refractory to medical treatment.
- 2) Turbinate reduction-Endoscopic sinus surgery is used for reduction of hypertrophied turbinates causing nasal obstruction.
- 3) Septoplasty-Deviated nasal septum causing nasal obstruction can be corrected by endoscopic sinus surgery.
- 4) Control of epistaxis-Endoscopic sinus surgery is useful in identification and ligation of blood vessels responsible for epistaxis.
- 5) Orbital decompression-Orbital decompression in patients with exophthalmos can be performed using endoscopic sinus surgery.
- 6) Repair of CSF leak-Endoscopic sinus surgery is used in repair of CSF leaks occurring through the anterior skull base.

- 7) Pituitary surgery-Endoscopic sinus surgery is used in performing transsphenoidal hypophysectomy.

### **COMPLICATIONS OF ENDOSCOPIC SINUS SURGERY**

1. Injury to nasolacrimal duct can lead to excessive tearing in the post operative period.
2. Injury to medial wall of orbit can occur leading to prolapse of orbital fat and orbital haemorrhage.
3. Injury to cribriform plate of ethmoid can lead to leakage of CSF
4. Surgeries involving sphenoid sinus can lead to injury to carotid artery and optic nerve.
5. Surgeries involving frontal sinus can also result in leakage of cerebrospinal fluid.

## **ANESTHETIC CONSIDERATIONS IN ENDOSCOPIC SINUS SURGERY**

### **PRE OP PREPARATION**

Patients undergoing endoscopic sinus surgery are premedicated with nasal decongestant drops like xylometazoline to reduce bleeding. Steroids like dexamethasone are given preoperatively to reduce nasal edema in the postoperative period.

### **INTRAOPERATIVE MANAGEMENT**

Endoscopic sinus surgery can either be performed under local anesthesia or monitored anesthesia care or general anesthesia. In patients with extensive disease, involvement of multiple sinuses and in patients with reactive airway disease surgery is preferably performed under general anaesthesia. Laryngeal mask airway can be used instead of an endotracheal tube as it prevents entry of blood into the stomach. Laryngeal mask airway is particularly useful in patients with reactive airway disease as it prevents coughing or bucking during emergence, which can lead to bleeding from the surgical site.

Maintenance of a bloodless field is essential during sinus surgery to improve visualization of the surgical field. It can be achieved by placing pledgets soaked with vasoconstrictive agents like adrenaline or xylometazoline in the nasal cavity or by blocking the sphenopalatine




ganglion using infiltration with 1% lignocaine and 1:1,00,000 epinephrine or by inducing hypotension with hypotensive agents.

## **POSTOPERATIVE MANAGEMENT**

Patients undergoing endoscopic sinus surgery should be monitored in the post op period as they are at the risk of aspiration of the swallowed blood. Pain relief should be given in the post op period using opioids.

## PHARMACOLOGY OF ESMOLOL

Esmolol is a cardioselective ultra short acting beta blocker. The chemical name of esmolol is methyl p-[2-hydroxy-3-(isopropylamino)propoxy] hydrocinnamate hydrochloride.





## **CLINICAL PHARMACOLOGY OF ESMOLOL**

Esmolol is a short acting agent with a elimination half life of 9 minutes .Though esmolol is a cardioselective beta blocker at higher doses it acts on beta 2 receptors present in bronchial smooth muscles and blood vessels.

### **PHARMACOKINETICS AND METABOLISM**

Esmolol is metabolized by RBC esterases into free acid and methanol. The acid metabolite is excreted through urine and hence in patients with limited renal reserve,the elimination half life of esmolol is prolonged. The rate of clearance of esmolol was 20L/kg/hr, which is greater than the cardiac output and it's metabolism is not affected by rate of blood flow to the liver and kidneys.

Esmolol is administered at a loading dose of 1 mg/kg and the infusion dose of esmolol is 0.05-0.3mg/kg/min. With loading dose steady state plasma concentration is achieved within 5 min's. With infusion steady state plasma concentration is achieved in 30 min.

### **PHARMACODYNAMICS**

Esmolol causes decrease in heart rate, delay in recovery of sinus node,Prolongation of AH interval during normal sinus rhythm and increase in antegrade wenckebach cycle length.

During exercise esmolol produces decrease in heart rate, blood pressure, rate pressure product, right and left ventricular ejection fraction and cardiac index. Discontinuation of esmolol is associated with return of parameters to baseline in about 30 minutes. Esmolol did not produce any adverse reactions in patients with bronchial asthma and COPD.

## **INDICATIONS**

- 1) Supraventricular tachycardia-Esmolol is used in rapid control of ventricular rate in patients with atrial fibrillation and atrial flutter.
- 2) Intraoperative and postoperative tachycardia and hypertension  
Esmolol is used to blunt the tachycardia and hypertension associated with laryngoscopy and intubation and during emergence from anesthesia.

For control of intraoperative tachycardia and hypertension esmolol is administered at a loading dose of 1mg/kg over one minute followed by infusion at a rate of 150microgram/kg/min.

## **CONTRAINDICATIONS**

- 1) Heart block
- 2) Sinus bradycardia
- 3) Congestive cardiac failure
- 4) Cardiogenic shock

## **ADVERSE EFFECTS**

- **Cardiovascular system**

Esmolol can produce hypotension, bradycardia, chest pain pulmonary edema, peripheral ischemia, syncope, heart block and even asystole.

- **Central nervous system**

Headache, paraesthesia, dizziness, anxiety, depression, asthenia, somnolence, lightheadedness, abnormal thinking, agitation and fatigue.

- **Respiratory system**

Bronchospasm, dyspnoea, wheezing, rhonchi and nasal congestion can occur.

- **Gastrointestinal system**

Dyspepsia, nausea and vomiting, constipation and abdominal discomfort can occur.

- **Skin**

Discoloration, edema, erythema, thrombophlebitis and skin necrosis can occur.

- **Miscellaneous**

Retention of urine, visual disturbance, speech disorder, fever and rigors can occur.

## **DRUG INTERACTIONS**

- 1) Administration of esmolol with morphine resulted in increase in plasma concentrations of esmolol.
- 2) Pharmacokinetics of esmolol was not affected by concomitant administration of digoxin and warfarin.
- 3) Concomitant administration of esmolol and reserpine resulted in worsening of bradycardia and hypotension.
- 4) Esmolol can cause prolongation of neuromuscular blockade caused by administration of succinyl choline.
- 5) Esmolol should be cautiously used in patients on calcium channel blockers like verapamil as it lead to heart block and cardiac arrest.
- 6) Esmolol should be cautiously used along with vasopressors like dopamine, dobutamine and noradrenalin as myocardial contractility can be impaired.
- 7) Esmolol should be cautiously used in patients who are already on drugs which impair myocardial contractility.
- 8) Patients on esmolol require a higher dose of epinephrine to treat anaphylactic reactions.

## **TOXICITY**

Esmolol overdose can cause cardiac arrest. Use of premixed injections can avoid the dilutional errors occurring with esmolol.

## **Management**

- 1) Hypotension should be treated with intravenous fluids and vasopressors.
- 2) Bradycardia should be treated with intravenous atropine.
- 3) Bronchospasm should be treated with beta2 agonists.
- 4) Cardiogenic shock should be treated with inotropic agents like dopamine, dobutamine, isoprenaline or using inodilators like milrinone.

## **PRECAUTIONS**


- 1) Esmolol should be cautiously used in diabetic patients as it can mask the symptoms of hypoglycemia.
- 2) Esmolol should be avoided in patients with end stage renal disease as the plasma concentrations of esmolol can get elevated in patients with renal disease.
- 3) Esmolol should be cautiously used in patients with bronchial asthma as bronchospasm can occur.



- 4) Esmolol should be avoided in pregnancy as fetal bradycardia can occur.
- 5) Esmolol should be avoided in nursing mothers as esmolol is excreted in breast milk.

## PHARMACOLOGY OF DEXMEDETOMIDINE

Dexmedetomidine is a selective  $\alpha$ -2 agonist which has sedative and analgesic properties.





## **PHARMACODYNAMICS**

Dexmedetomidine is selective for alpha 2 receptors but at higher doses it also has alpha 1 action. The sedative properties of dexmedetomidine are mediated through action on post synaptic alpha 2 receptors in locus caeruleus in brain and spinal cord. Dexmedetomidine has also action on muscarinic, dopaminergic, adrenergic and serotonin receptors.

## **PHARMACOKINETICS**

Dexmedetomidine has a distribution half life of about 6 minutes and a elimination half life of about 2 hours. The volume of distribution is 118 litres and it is highly protein bound .

Metabolism-Dexmedetomidine is predominantly metabolized in the liver into inactive metabolites. Metabolism occurs through glucuronidation and through cytochrome p450 pathways. Context sensitive half life of dexmedetomidine is 4 to 10 minutes.

### **Elimination**

Elimination of dexmedetomidine takes place through the kidneys and hence dexmedetomidine should be cautiously used in patients with severe renal impairment.

## **INDICATIONS**

1. Dexmedetomidine is principally used for sedation of mechanically ventilated patients in the ICU. Dexmedetomidine has the advantage of producing sedation without significant respiratory depression.
2. Dexmedetomidine can also be used for sedation during surgical procedures in monitored anaesthesia care.
3. Dexmedetomidine can be used for rapid opioid detoxification, cocaine withdrawal and intolerance after prolonged use of benzodiazepines and opioids.
4. Dexmedetomidine can also be used in awake fiber optic intubation, awake craniotomies with functional testing and electrocorticography and in awake carotid endarterectomies with stable haemodynamics.

## **CONTRAINDICATIONS**

1. Contraindicated in patients with severe hepatic and renal dysfunction.
2. Patients with known hypersensitivity to dexmedetomidine.

## **DOSE**

Dexmedetomidine is administered at a loading dose of 1 microgram/kg followed by infusion at a rate of 0.5 micrograms/kg/hour.

Infusion dose can be increased by 0.1 microgram/ kg/hour every 5 minutes if adequate clinical response is not obtained. Elderly patients are highly sensitive to dexmedetomidine, Hence the dose of dexmedetomidine should be reduced. Effects of dexmedetomidine can be antagonized by administration of atipamazole.

## **ADVERSE EFFECTS**

The common adverse effects of dexmedetomidine are hypotension, bradycardia, dryness of mouth, hypertension and nausea.

Other adverse effects include fever, arrhythmias like atrial fibrillation, edema, myocardial infarction, pulmonary edema, speech disorders, diarrhoea, hyperkalemia, hyperglycemia, muscle weakness, paraesthesia, delirium, hallucinations, depression, urinary retention, hypoxia, hypercapnia, hypoventilation, pulmonary hypertension, pneumothorax, erythematous rashes and visual disturbances.

Administration of dexmedetomidine is associated with elevation of liver enzymes and decrease in prothrombin levels. Dexmedetomidine can produce withdrawal syndrome after discontinuation.

## **DRUG INTERACTIONS**

Dexmedetomidine enhances the effects of anaesthetic agents such as hypnotics, sedatives and opioids. Hence, co-administration of these drugs with dexmedetomidine results in decrease in dose requirement of

these agents. Caution should be exercised during administration of dexmedetomidine along with vasodilators and negative inotropic agents as it can worsen bradycardia and hypotension associated with dexmedetomidine. Dexmedetomidine should be avoided in pregnancy and during labour and caesarean section.

Dexmedetomidine should be avoided in nursing mothers as it is excreted in breast milk.

## **PRECAUTIONS**

Close monitoring of the patient with ECG, Pulse oxymeter and Blood pressure monitoring should be done during administration of dexmedetomidine.

Caution should be exercised while administering dexmedetomidine to patients with pre-existing heart block, Congestive cardiac failure and bradycardia.

Patient should be adequately hydrated prior to administration of dexmedetomidine to prevent hypotension.

Treatment of hypotension includes stoppage of the infusion, elevation of the foot end of the patient and administration of vasopressors.

Administration of loading dose of dexmedetomidine can be associated with transient hypertension. This occurs due to action on alpha 1 receptors in blood vessels due to the high plasma concentration attained during administration of the loading dose.

Rapid intravenous administration of dexmedetomidine to individuals with high vagal tone can cause severe bradycardia and cardiac arrest. Treatment of bradycardia involves administration of anticholinergic agents such as atropine or glycopyrrolate.

Administration of dexmedetomidine can cause dryness of eyes and hence patient's eyes should be lubricated to prevent corneal dryness.

Sudden discontinuation of dexmedetomidine after prolonged infusion can cause withdrawal syndrome characterized by rebound hypertension, headache, agitation and nervousness. Hence infusion of dexmedetomidine should be limited to less than 24 hours.



## REVIEW OF LITERATURE

**Boezaart et al** (1995) in a prospective study of forty patients to compare the effects of sodium nitroprusside and esmolol in controlled hypotension for functional endoscopic sinus surgery. They found that in esmolol group ,due to unopposed alpha adrenergic effect on mucous membrane superior surgical conditions were obtained with mild hypotension(MAP>65 mmhg) when compared with nitroprusside group.

**Koi io et al**(2009) in a prospective double blinded study of 48 patients to compare the effects of combining desflurane with esmolol and dexmedetomidine in controlled hypotension for tympanoplasty surgery found that both esmolol and dexmedetomidine can be effectively used along with desflurane for tympanoplasty surgery and esmolol group had shorter recovery time and less post operative sedation when compared with dexmedetomidine group.

**Turan et al**(2007) in a prospective randomized study of patients to compare the effects of dexmedetomidine, remifentanil and esmolol in controlled hypotensive anesthesia found that no significant differences were seen between threegroups with respect to quality of surgical field and intra operative MAP and heart rate but post extubation values were

significantly lower in remifentanyl and esmolol group when compared with dexmedetomidine group.

**Baysoni et al** (2013) in prospective randomized study for comparison of remifentanyl and esmolol in induced hypotension for nasal surgery in 30 patients found no significant differences between the two groups with respect to intra operative surgical field and vital parameters like MAP and heart rate and concluded that remifentanyl and esmolol provided dry surgical fields during nasal surgery.

**Y j lim et al**(2003) in a study to compare esmolol with esmolol and normovolemic haemodilution for controlled hypotension in 30 patients undergoing spinal surgery found that transfusion of blood products in esmolol normovolemic haemodilution group was significantly lower when compared with esmolol alone group and concluded that esmolol in combination with normovolemic haemodilution can be effectively used for blood conservation in spinal surgeries.

**Celebi et al**(2007) in a prospective study of comparison of esmolol and remifentanyl on cognitive functions during induced hypotension for tympanoplasty surgeries found that both remifentanyl and esmolol provided ideal surgical conditions and they produced

similar effects on cognitive function in the immediate post operative period.

**Guney et al**(2002) in a prospective study of esmolol compared with nitroglycerine for controlled hypotension in patients undergoing nasal surgeries found that ideal surgical conditions and better haemodynamic stability was provided with esmolol than nitroglycerine. They concluded that esmolol can be effectively used as alternative to nitroglycerine for controlled hypotension.

**Eun et al**(1997) in a comparative study between sodium nitroprusside and esmolol with induced hypotension for total hip arthroplasty in 70 patients found there were no significant differences between the two groups with respect to blood loss, blood transfusion and fluid administration and concluded that esmolol can be effectively used as an hypotensive agent in hip arthroplasties without significant side effects.

**Gokce et al** (2009) in prospective randomized double blinded study with 40 patients of hypotensive anesthesia with esmolol for assessment of haemodynamics and consumption of anesthetic agents and recovery studied the effects of adding esmolol to propofol and remifentanyl combination in controlled hypotension and concluded that

the addition reduced the requirement of remifentanyl without decrease in cardiac output.

**Kumsuk park et al(2006)** in a prospective study of hydralazine pretreatment in patients undergoing controlled hypotensive anesthesia with esmolol in spine surgery found that time required to achieve the target mean arterial pressure was significantly lowered in the group, which had hydralazine pretreatment than esmolol alone group without any decrease in cardiac output and concluded that addition of hydralazine increases the efficacy of esmolol and decreases its dose requirement without decrease in cardiac output.

**Himanshudongre et al(2012)** in a study of controlled hypotensive anesthesia comparing esmolol and nitroglycerine in producing a dry operating field for spinal surgery found that the heart rate and mean arterial pressure required for ideal surgical field were significantly lower in esmolol group than nitroglycerine group.

**Erbesler et al (2013)** in a prospective study compared the effectiveness of dexmedetomidine and esmolol in induced hypotension on clinical course and cost and found that the recovery index and duration of clinical action were significantly prolonged in dexmedetomidine group than esmolol group and concluded that both

drugs can be used in induced hypotension but neuromuscular blocking time was prolonged in dexmedetomidine group.

**Mengistu et al** (2007) studied the effects of sodium nitroprusside and esmolol used in induced hypotension on natriuretic peptides in 45 patients undergoing endoscopic nasal surgery. Patients were divided into three groups (control, esmolol, nitroprusside group), they found that levels of ANP (atrial natriuretic peptide) and BNP (brain natriuretic peptide) were significantly lower in nitroprusside and esmolol group than control group and concluded that decrease in natriuretic peptide levels were due to decrease in cardiac filling pressures due to decrease in systemic vascular resistance associated with hypotensive anaesthesia.

**Zhou qi et al** (2010) in a prospective study evaluated the effects of Combining esmolol with nitroglycerine for induced hypotension in Hypertensive patients undergoing spinal surgery and concluded that the addition of esmolol with nitroglycerine has decreased blood loss in the intraoperative period and the duration of surgery.

**Akkaya et al** (2014) in a randomized clinical study compared the effects of dexmedetomidine and magnesium sulphate on the quality of surgical field in patients undergoing endoscopic sinus surgery. They found that heart rate and duration of surgery were significantly lower in

dexmedetomidine group, and the quality of surgical field was much better in dexmedetomidine group than magnesium sulphate group. They concluded that dexmedetomidine was a much better alternative to magnesium sulphate for induced hypotension.

**Shen et al** (2001) in a double blinded randomized control trail, Studied the effects of intra operative esmolol infusion on quality of surgical fields during sevoflurane anesthesia in 40 patients undergoing endoscopic sinus surgery. The patients were divided into two groups ,one received esmolol infusion and other received saline infusion along with inhalational agent sevoflurane. They found that heart rate and blood pressure were significantly lower in esmolol group and the quality of surgical field was better in esmolol group. They also found that blood loss was much lower in esmolol group. They concluded that intra operative infusion of esmolol produced bradycardia and hypotension and the quality of surgical field was significantly improved in patients undergoing endoscopic sinus surgery.

**Abdulla aydin ozcan et al** (2012) in a randomized prospective study Compared remifentanyl and dexmedetomidine for induced hypotension in 50 patients undergoing endoscopic sinus surgery. They found that heart rate was higher in remifentanyl group than

dexmedetomidine group and the recovery time was prolonged in dexmedetomidine group than remifentanyl group. They concluded that both remifentanyl and dexmedetomidine can be effectively used for producing improved quality of surgical field in patients undergoing endoscopic sinus surgery but dexmedetomidine was associated with prolonged recovery time than remifentanyl.

**Farah nasreen et al**(2009) studied the effects of using dexmedetomidine for controlled hypotension in 42 patients undergoing middle ear surgery. The patients were divided into two groups, one received infusion of dexmedetomidine and other group received saline infusion. The target mean arterial pressure was achieved with halothane. They found that the requirement of halothane to achieve the target mean arterial pressure was much lower in dexmedetomidine group and surgical field quality was much better in dexmedetomidine group. They concluded that hypotensive anesthesia can be safely provided in patients undergoing middle ear surgery with dexmedetomidine.

**Richaf et al** (2008) in a prospective study compared dexmedetomidine with remifentanyl in controlled hypotension for patients undergoing tympanoplasty. They concluded that

dexmedetomidine can be safely used as an alternative to remifentanyl for providing hypotensive anesthesia in tympanoplasty surgery.

**Bayazit dikmen et al**(2010)studied the effects of using dexmedetomidine with low flow anesthesia for controlled hypotension in 40 patients undergoing middle ear surgery.The patients were divided into two groups, one received dexmedetomidine infusion and another group received saline infusion. They found that the dexmedetomidine group achieved target mean arterial pressure earlier than saline group.They concluded that dexmedetomidine effectively produced sustained hypotension with low flow anesthesia in patients undergoing middle ear microsurgery.

**Malhotra et al**(2013)in a prospective study,studied the effects of dexmedetomidine in improving the quality of surgical field and the requirement of hypotensive agents in 60 patients undergoing endoscopic sinus surgery under general anaesthesia. Patients were divided into two groups,one received dexmedetomidine infusion and another received saline infusion and found that nitroglycerine needed to achieve target mean arterial pressure was lower in dexmedetomidine group and mean arterial pressure was significantly lower in dexmedetomidine group. They concluded that better quality surgical



field can be provided with hypotensive anesthesia using dexmedetomidine.

**Guang jiegao et al**(2012)studied the feasibility of dexmedetomidine for induced hypotension in 48 patients undergoing endoscopic sinus surgery with sevoflurane. The patients were divided into two groups, dexmedetomidine group and saline group. They found that the dose of anesthetic agents like propofol, sevoflurane and fentanyl were significantly lower in dexmedetomidine group ,mean arterial pressure and heart rate were also significantly lower in dexmedetomidine group, incidence of shivering and delirium were also lower in dexmedetomidine group but the incidence of bradycardia was higher in dexmedetomidine group. They concluded that dexmedetomidine provided cardiovascular stability and reduced the intraoperative requirement of anesthetic agents and was effective for hypotensive anesthesia in endoscopic sinus surgery.

**Sunil chiruvella et al**(2014)made a prospective randomized double blinded study with dexmedetomidine for controlled hypotensive anesthesia in endoscopic sinus surgery.They found that duration of surgery and mean arterial pressure were lower in dexmedetomidine group and the quality of surgical field was better in dexmedetomidine

group. They concluded that dexmedetomidine can be safely used for induced hypotension during endoscopic sinus surgery.

**Gurbet et al** (2006) studied the effects of intraoperative infusion of dexmedetomidine on perioperative analgesic requirements in 50 patients. The patients were divided into two groups, one group received dexmedetomidine infusion and another group received saline infusion.

They found that requirement of morphine in the post operative period was significantly lower in dexmedetomidine group. They concluded that intra operative use of dexmedetomidine provides effective post operative analgesia and reduces analgesic requirement in the post operative period without increasing side effects.

**Kim sy et al**(2013) studied the effects of intraoperative use of dexmedetomidine infusion on agitation during emergence and recovery quality after nasal surgery in 100 patients. Patients were divided into two groups, one received dexmedetomidine infusion and another received saline infusion. They found that emergence agitation was lower in dexmedetomidine group and heart rate and blood pressure were lower in dexmedetomidine group. 24 hours after surgery global –QOR-40 score was higher in dexmedetomidine group. They concluded that intra

operative infusion of dexmedetomidine resulted in better recovery and smooth emergence after nasal surgery.

**Guray demir et al**(2009) studied the effects of dexmedetomidine on hemodynamic parameters and dose of propofol during anesthetic induction in 80 patients. Patients were divided into two groups, one group received dexmedetomidine and another group received saline. They found that mean arterial pressure and heart rate were significantly lower in dexmedetomidine group, the dose of propofol was also lower in dexmedetomidine group. They concluded the use of propofol before induction blunted hemodynamic response associated with laryngoscopy and intubation, it also reduced the requirement of propofol.

## **MATERIALS & METHODS**

The study was a prospective randomized single blinded control trial involving 50 patients (25 in each group) attending the Department of Ear, nose and Throat, Chengalpattu Medical College Hospital over a period of one year(July 2013 – June 2014). Institutional ethical committee approval was obtained. The procedure was explained to the patient. Informed consent was obtained. Detailed history of the patient was collected. Routine investigations like complete blood count, blood glucose, renal function tests, serum electrolytes, chest x ray, electrocardiogram were done as per our hospital protocol. Patients fulfilling the inclusion criteria and who gave consent were then randomly allocated to one of the study groups on the basis of computerized randomized list.

### **Inclusion Criteria:**

1. ASA 1
2. Age: 20-50 years
3. Patients posted for Elective FESS

## **Exclusion Criteria**

1. Patients with coronary artery disease
2. Patients with diabetes
3. Patients with Hypertension
4. Patients with renal dysfunction
5. Patients with Hepatic Dysfunction
6. Patients with Bleeding Diathesis,
7. Patients with Cerebrovascular disease
8. Patients with Recurrent sinus surgery

## **Materials Needed**

1. 18G iv Cannula
2. 18G radial artery cannula
3. 50cc Syringe
4. Infusion Pump
5. Transducer
6. 2cc, 5cc and 10cc Syringes
7. Injection Dexmedetomidine

8. Injection Esmolol
9. Boyle's Apparatus
10. Working laryngoscope, cuffed endotracheal tubes of appropriate size, Airway, suction apparatus with suction catheter.
11. Emergency drugs like Inj. Adrenaline, Inj. Atropine, Inj. Ephedrine.
12. Monitor for continuous monitoring of Invasive blood pressure, ECG, Respiratory rate, Oxygen saturation.

### **Methodology**

18 G iv Cannula was inserted and the patients started on infusion Ringer Lactate. Another 18G iv Cannula was inserted in the opposite limb for infusion of the Study drug.

18 G radial Artery Cannula was inserted for continuous measurement of Arterial Blood Pressure.

All patients were premedicated with Injection Glycopyrolate 0.2mg intramuscularly 45 minutes prior to surgery, Injection Midazolam 0.05 mg/kg intravenously and Injection Fentanyl 2 microgram/kg intravenously.

In Esmolol group, patients were given a loading dose of Esmolol at 1mg/kg over 1 minute followed by infusion at the rate of 0.4-0.8 mg/kg/hour before induction. The infusion dose was titrated to obtain a mean arterial pressure between 55-65 mm Hg.

In Dexmedetomidine group, patients were given a loading dose Of dexmedetomidine at 1 microgram/kg diluted in 10 mL of 0.9 % Saline over 10 minutes followed by infusion at the rate 0.4-0.8 micrograms/kg/hour before induction. The infusion dose was titrated to obtain a mean arterial pressure between 55-65 mm Hg.

Induction agent used in all patients in this study was Propofol. Propofol was administered at a dose of 1-2mg/kg until the loss of Verbal response. Induction dose of Propofol was recorded.

Intubation was done following administration of Injection Atracurium 0.5 mg/kg with appropriate sized Endotracheal Tubes. Anaesthesia was maintained with 60 % N<sub>2</sub>O/O<sub>2</sub> mixture.

All patients were placed in 15 degree reverse Trendelenberg position and their nasal cavities were packed with Cottonoids, soaked with Epinephrine in a concentration of 1: 80,000.

All the surgeries were performed by the same Surgeon, who was blinded to the hypotensive agent used, so as to ensure consistency in estimating the quality of the surgical field.

Infusion of the study drugs was stopped 10 minutes before the end of the surgery. Patients were extubated after reversal with inj neostigmine and inj glycopyrolate.

### **Parameters Monitored**

#### **1. Dose of Propofol**

2. **Blood Pressure**- Pre-operatively, After induction, at 15 minutes, at 30 minutes, at 45 minutes during surgery, 5 minutes after stoppage of infusion of study drug, 10 minutes after stoppage of infusion of study drug, at the end of surgery, after extubation.

3. **Heart rate** - Pre-operatively, After induction, at 15 minutes, at 30 minutes, at 45 minutes during surgery, 5 minutes after stoppage of infusion of study drug, 10 minutes after stoppage of infusion of study drug, at the end of surgery, after extubation.



4. **Serum cortisol level intraoperatively**

5. **Emergence Time:** It is the time interval between stoppage of the anaesthetic drugs and eye-opening to oral commands.

6. **Quality of the surgical field-** It is assessed using average category scale.

**Average Category Scale**

Score 0	-	No bleeding
Score 1	-	Minimal Bleeding, suctioning not required
Score 2	-	Minimal Bleeding, suctioning occasionally required
Score 3	-	Minimal Bleeding, repeated suctioning required
Score 4	-	Moderate Bleeding, repeated suctioning required, bleeding obscures surgical field
Score 5	-	Severe Bleeding, surgery not possible as bleeding completely obscures the surgical field

7. **PostAnaesthesia Recovery Score:** it is assessed using **Modified Aldrete Score.**

1. **Activity-** Able to move voluntarily or on command

Four extremities	2
------------------	---

Two extremities	1
-----------------	---

No extremities 0

**2. Respiration**

Able to breathe deeply and cough freely 2

Dyspnea, shallow or limited breathing 1

Apnea 0

**3. Circulation**

Blood pressure within 20 mm Hg of Pre-Op level 2

Blood Pressure within 20-50 mm Hg of Pre-Op level 1

Blood Pressure  $\pm$  50 mm Hg of Pre-Op level 0

**4. Consciousness**

Fully Awake 2

Arousable on command 1

unresponsive 0

**5. Oxygen saturation**

Saturation more than 92 % 2

Needs Oxygen to maintain saturation more than 90 % 1

Saturation less than 90 % with oxygen 0

Total Score- 15

9 or more points are required for recovery to be confirmed.

Time required to attain Modified Aldrete Score of more than 9 is recorded.

8. **Sedation Score** at 15, 30 and 60 minutes after extubation is recorded.

**Sedation Score- Ramsay Sedation Score**

Score 1 Patient is anxious, agitated or restless or both

Score 2 Patient is cooperative, oriented and tranquil

Score 3 Patient responds to commands only

Score 4 Patient exhibits brisk response to light, glabellar tap or loud auditory stimulus

Score 5 Patient exhibits sluggish response to light, glabellar tap or loud auditory stimulus

Score 6 Patient exhibits no response

9. The time required for first rescue analgesic in the post-operative period is also recorded.

## **STATISTICAL ANALYSIS**

Dose of propofol, heart rate, systolic blood pressure, diastolic blood pressure, emergence time, post anaesthesia recovery score, sedation score, time for rescue analgesia. All recorded data were entered using MS-excel software and analysed using SPSS(Statistical Package of Social Sciences, Version 16 for windows, student's t-test was used to compare the two groups on mean values of various parameters. The p-value taken for significance is  $<0.05$ .

## **RESULTS AND OBSERVATION**

Our study included 50 patients, patients were randomly allocated into either group E (esmolol group) or group D (dexmetetomidine). Initial patient characteristics such as age, weight and ASA grading were noted.


Outcomes measured are dose of propofol, heart rate and blood pressure, quality of surgical field, emergence time, post anesthesia recovery score, sedation score at 15 minutes, 30 minutes and 60 minutes after surgery, time for rescue analgesia post operatively, serum cortisol intraoperatively.

## PATIENT CHARECTERISTICS

### AGE(student's t test)

The average age did not differ between the two groups. The mean age in group E was  $31.84 \pm 8.68$  years and that of group D was  $33.08 \pm 7.39$ . The difference was not statistically significant(p value =0.5).


Group	Mean	T value	P value
Group D	$33.08 \pm 7.3$	0.543	0.5
Group E	$31.84 \pm 8.6$		



### WEIGHT(student's t test)

The average weight did not differ between the two groups. The average weight of patients in group E was  $57.04 \pm 9.74$  and the average weight of patients in group D was  $56 \pm 9.12$ . The difference was not statistically significant (p value = 0.7).

Group	mean	T value	P value
Group D	$56 \pm 9.1$	0.389	0.7
Group E	$57.04 \pm 9.7$		




## OUTCOMES MEASURED

### 1) Dose of propofol(student's t test)

The induction dose of propofol was lower in group D( $1.24 \pm 0.10$  mg/kg) when compared with group E( $1.65 \pm 0.13$  mg/kg), which was statistically significant ( $p$  value  $< 0.001$ ).

Group	mean	T value	P value
Group D	$1.24 \pm 0.10$	12.20	0.00
Group E	$1.65 \pm 0.13$		







## 2) HEART RATE(student's t test)

Heart rates were compared between two groups preoperatively, after induction, at (15 minutes, 30 minutes, 45 minutes) during surgery, at (5 minutes and 10 minutes after stoppage of study drug), at end of surgery, after extubation.


Comparison showed no statistically significant differences in heart rate between two groups preoperatively (p value=0.058)




Comparison showed no statistically significant differences in heart rate between two groups after intubation(p value=0.27).




Comparison showed no statistically significant differences in heart rate between two groups at 15 minutes during surgery (pvalue=0.082)




Comparison showed no statistically significant differences in heart rate between two groups at 30 minutes during surgery(pvalue=0.05)



Comparison showed no statistically significant differences in heart rate between two groups at 45 minutes during surgery( p value=0.09).




Comparison showed statistically significant differences in heart rate between two groups at 5 minutes after stoppage of study drug ( $p$  value $<0.001$ ). Heart rate's were significantly higher in group E than group D.




Comparison showed statistically significant differences in heart rate between two groups at 10 minutes after stoppage of study drug


(p value=0.002).Heart rate's were significantly higher in group E than group D.




Comparison showed statistically significant differences in heart rate between two groups at end of surgery(p value=0.002).Heart rate's were significantly higher in group E than group D.



Comparison showed statistically significant differences in heart rate between two groups at end of surgery(p value<0.001).Heart rate's were significantly higher in group E than group D.



<b>Time</b>	<b>Group</b>	<b>Mean</b>	<b>T value</b>	<b>p value</b>
Pre operative	D	83.04±8.1	1.944	0.058
	E	87.52±8.1		
After intubation	D	73.6±6.8	1.117	0.27
	E	71.32±7.5		
15 min of surgery	D	67.12±5.7	1.777	0.082
	E	64±6.6		
30 min of surgery	D	64.44±8.5	2.34	0.05
	E	62.56±6.1		
45 min of surgery	D	62.52±6.0	1.732	0.09
	E	54.44±6.5		
5 min after stoppage	D	60.44±6.7	3	0.00
	E	66.08±6.5		
10 min after stoppage	D	65.12±5.8	3.284	0.002
	E	70.96±6.6		
At end of surgery	D	67.27±5.7	3.351	0.002
	E	73.12±6.6		
After extubation	D	68.76±5.3	7.576	0.00
	E	81.12±6.1		







### 3. COMPARISON OF SYSTOLIC BLOOD PRESSURE(student's t test)

Systolic blood pressure was compared between two groups preoperatively,after induction, at(15 minutes,30 minutes,45 minutes ) during surgery,at(5 minutes and 10 minutes after stoppage of study drug), at end of surgery, after extubation.


Comparison showed no statistically significant differences in systolic blood pressure between two groups pre operatively (p value=0.98)




Comparison showed no statistically significant differences in systolic blood pressure between two groups after intubation (p value=0.06).




Comparison showed no statistically significant differences in systolic blood pressure between two groups at 15 minutes during surgery (p value=0.05).




Comparison showed no statistically significant differences in systolic blood pressure between two groups at 30 minutes during surgery (p value=0.08).




Comparison showed no statistically significant differences in systolic blood pressure between two groups at 45 minutes during surgery (p value=0.17).




Comparison showed statistically significant differences in systolic blood pressure between two groups at 5 minutes after stoppage of study drug. Systolic blood pressure was significantly higher in group E than group G (p value<0.001).




Comparison showed statistically significant differences in systolic blood pressure between two groups at 10 minutes after stoppage of study drug. Systolic blood pressure was significantly higher in group E than group G (p value<0.001).




Comparison showed statistically significant differences in systolic blood pressure between two groups at end of surgery. Systolic blood pressure was significantly higher in group E than group G ( $p$  value  $< 0.001$ ).



Comparison showed statistically significant differences in systolic blood pressure between two groups after extubation. Systolic blood pressure was significantly higher in group E than group G (p value<0.001).



<b>Time</b>	<b>Group</b>	<b>Mean</b>	<b>T value</b>	<b>P value</b>
Pre operatively	D	123.24±10.8	0.03	0.98
	E	123.32±7.6		
After intubation	D	110.72±5.5	1.92	0.06
	E	114.32±6.7		
15 min of surgery	D	91.72±4.9	1.91	0.05
	E	93.08±5.5		
30 min of surgery	D	88.36±3.5	1.813	0.08
	E	90.28±3.8		
45 min of surgery	D	89.6±2.9	1.407	0.17
	E	90.92±3.6		
5 min after stoppage	D	92.48±2.6	14.54	0.00
	E	112.16±6.1		
10 min after stoppage	D	94.32±2.8	16.69	0.00
	E	115.32±5.6		
At end of surgery	D	95.32±3.0	15.65	0.00
	E	116.4±5.4		
After extubation	D	101.2±3.7	11.53	0.00
	E	118.24±6.3		







## COMPARISON OF DIASTOLIC BLOOD PRESSURE(student's t test)

Diastolic blood pressure was compared between two groups preoperatively, after induction, at(15 minutes,30 minutes,45 minutes) during surgery, at(5 minutes and 10 minutes after stoppage of study drug), at end of surgery, after extubation.


Comparison showed no statistically significant differences in diastolic blood pressure between two groups pre operatively (p value=0.27).




Comparison showed no statistically significant differences in diastolic blood pressure between two groups after intubation (p value=0.31).




Comparison showed no statistically significant differences in diastolic blood pressure between two groups at 15 minutes during surgery (p value=0.54).




Comparison showed no statistically significant differences in diastolic blood pressure between two groups at 30 minutes during surgery (p value=0.15).




Comparison showed no statistically significant differences in diastolic blood pressure between two groups at 45 minutes during surgery (p value=0.19).




Comparison showed statistically significant differences in diastolic blood pressure between two groups at 5 minutes after stoppage of study drug .diastolic blood pressure was significantly higher in group E than group G (p value<0.001).




Comparison showed statistically significant differences in diastolic blood pressure between two groups at 10 minutes after stoppage of study drug .diastolic blood pressure was significantly higher in group E than group G (p value<0.001).




Comparison showed statistically significant differences in diastolic blood pressure between two groups at end of surgery .diastolic blood pressure was significantly higher in group E than group G (p value<0.001).



Comparison showed statistically significant differences in diastolic blood pressure between two groups after extubation .Diastolic blood pressure was significantly higher in group E than group G (p value<0.001).




<b>Time</b>	<b>Group</b>	<b>mean</b>	<b>T value</b>	<b>P value</b>
Pre op	D	76.36±6.5	1.118	0.27
	E	78.48±6.8		
After intubation	D	71.88±5.9	1.023	0.31
	E	73.56±5.6		
15 min of surgery	D	60.24±5.5	0.624	0.54
	E	61.48±8.2		
30 min of surgery	D	58.92±3.7	1.452	0.15
	E	60.52±4.0		
45 min of surgery	D	59.6±4.5	1.345	0.19
	E	61.16±3.5		
5 min after stoppage	D	61.28±4.1	7.624	0.00
	E	72.72±6.6		
10 min after stoppage	D	61.2±4.0	12.041	0.00
	E	77.4±5.3		
At end of surgery	D	64.48±3.3	12.604	0.00
	E	79.72±5.0		
After extubation	D	67.48±3.7	11.05	0.00
	E	80.08±4.3		



#### 4) QUALITY OF SURGICAL FIELD-AVERAGE CATEGORY

**SCALE(student's t test)**

Comparison of quality of surgical field showed no statistical difference between two groups.






##### 5) **EMERGENCE TIME(student's t test)**

Comparison of emergence time showed statistically significant differences between two groups. Emergence time was significantly higher in group D than group E (P value<0.001).


Group	Mean	T value	P value
Group D	13.14±1.1	15.57	0.00
Group E	7.62±1.3		



**6) POST ANESTHESIA RECOVERY SCORE-MODIFIED  
ALDRETE SCORE(student's t test)**

Comparison showed statistically significant differences between two groups in the time to achieve modified aldrete score of  $\geq 9$ . The time was significantly higher in group D compared with group E (P value  $<0.001$ ).

Group	Mean	T value	P value
Group D	15.4 $\pm$ 1.8	10.89	0.00
Group E	10.86 $\pm$ 0.9		




## 7) SEDATION SCORE-RAMSAY SEDATION

### SCORE(student's t test)

Comparison between two groups showed that there was statistically significant differences in sedation scores at 15 minutes (p value <0.001) and 30 minutes (p value=0.01) after surgery.

Sedation scores were significantly higher in group D than group E. There was no statistically significant difference in sedation score between two groups at 60 minutes after surgery.


Time	Group	mean	T value	P value
15 minutes	Group D	2.76±0.4	3.05	0.00
	Group E	2.36±0.4		
30 minutes	Group D	2.64±0.4	2.683	0.01
	Group E	2.28±0.4		
60 minutes	Group D	2		
	Group E	2		



## 8) RESCUE ANALGESIA(student's t test)

Comparison between two groups showed that there was statistically significant differences in the time for rescue analgesia (p value<0.001).The time for rescue analgesia was significantly higher in group D than group E.


Group	Mean	T value	P value
Group D	64.4±7.9	12.28	0.00
Group E	40.64±5.5		



**9) SERUM CORTISOL(student's t test)**

Comparison between two groups showed that there was no statistically significant difference in serum cortisol levels intraoperatively (p value=0.07).

Group	Mean	T value	P value
Group D	4.04±0.5	1.83	0.07
Group E	3.75±05		



## **DISCUSSION**

Functional endoscopic sinus surgery is one of the routinely performed surgeries. The use of hypotensive anesthesia during endoscopic sinus surgery has greatly reduced blood loss and improved visibility and quality of surgical field. Many studies have been published, which have stated the advantages of employing hypotensive anesthesia during endoscopic sinus surgery.

In our study we have compared the effects of dexmedetomidine given at a loading dose of 1 microgram/kg over 10 minutes followed by infusion at 0.4-0.8 microgram/kg /hr with esmolol given at a loading dose of 1mg/kg over 1 minute followed by infusion at 0.4-0.8 mg/kg/hr in induced hypotension for endoscopic sinus surgery.

The patients were comparable in both groups with regard's to age and weight.

### **DOSE OF PROPOFOL**

In our study, we found that the dose of propofol required for induction was significantly lower in dexmedetomidine group than esmolol group. The dose of propofol in dexmedetomidine group was (1.24±0.10 mg/kg) and in esmolol group it was (1.65±0.13 mg/kg).

This was consistent with the results obtained by

- 1) Suvadeep et al(2013) studied the effects of intraoperative dexmedetomidine infusion on requirement of propofol for maintainance of optimum depth of anesthesia during spine surgeries and concluded that dose of propofol was significantly reduced with the use of dexmedetomidine infusion.
- 2) Kanka et al(2009) studied the effects of dexmedetomidine on anaesthetic requirements in patients undergoing cardiovascular surgery and concluded that the dose of propofol and fentanyl were significantly lower with the use of dexmedetomidine.

## **HEART RATE AND BLOOD PRESSURE**

In our study, we found that the heart rate and blood pressure decreased significantly following administration of loading dose of esmolol and dexmedetomidine followed infusion heart rate and blood pressure intraoperatively (at 5 minutes and 10 minutes after stoppage of infusion) , at the end of surgery and after extubation were significantly lower in dexmedetomidine group than esmolol group.

This was consistent with the results obtained by



- 1) Shen et al (2001) studied the effects of intravenous infusion of esmolol on patients undergoing endoscopic sinus surgery and found that heart rate and blood pressure significantly decreased with esmolol infusion.
- 2) Malhotra et al (2013) studied the effects of dexmedetomidine on hypotensive anesthesia in patients undergoing hypotensive anesthesia and found that mean arterial pressure and heart rate was significantly lower with use of dexmedetomidine .

## **QUALITY OF SURGICAL FIELD**

In our study, the quality of surgical field was assessed using average category scale and we found that both esmolol and dexmedetomidine were effective in producing a surgical field with improved visibility (average category scale=2)

This is consistent with the results produced by

- 1) Farah nasreen et al (2009) studied the effects of dexmedetomidine for patients undergoing middle ear surgery with hypotensive anesthesia and found that the quality of surgical field was greatly improved with use of dexmedetomidine.

2) Boezaart et al(1995) made a comparative study between sodium nitroprusside and esmolol using inducing hypotension in functional endoscopic sinus surgery and found that superior surgical conditions were seen with esmolol even with mild hypotension(MAP>65 mm hg).

### **EMERGENCE TIME**

In our study, we found that the emergence time was significantly prolonged in dexmedetomidine group ( $13.14 \pm 1.13$  minutes) than esmolol group ( $7.62 \pm 1.36$  minutes).

This was consistent with the results obtained by

1) Abdulla aydin ozcan et al(2012) made a comparative study between remifentanyl and dexmedetomidine for controlled hypotension in patients undergoing endoscopic sinus surgery and found that recovery time was prolonged in dexmedetomidine group than remifentanyl group.

### **POST ANESTHESIA RECOVERY SCORE**

In our study post anesthesia recovery score was assessed using modified aldrete score .we found that post anesthesia recovery (time to achieve modified aldrete score of  $\geq 9$ ) was significantly higher in

dexmedetomidine group( $15.4 \pm 1.84$  minutes) than esmolol group ( $10.86 \pm 0.97$  minutes).

This was consistent with the results obtained by

- 1) Koi io et al(2009) made a comparative study between esmolol and dexmedetomidine combined with desflurane for controlled hypotension during tympanoplasty in adults and found that esmolol group had shorter recovery time than dexmedetomidine group.
- 2) Turan et al(2007) made a comparative study between esmolol, remifentanyl and dexmedetomidine in controlled hypotensive anesthesia and found that post extubation recovery score was longer in dexmedetomidine group than esmolol and remifentanyl group.

## **SEDATION SCORE**

In our study post operative sedation scores were assessed using Ramsay sedation score. we found that sedation scores were higher in dexmedetomidine group than esmolol group at 15 minutes( $2.76 \pm 0.43$  vs  $2.36 \pm 0.49$ ) and 30 minutes( $2.64 \pm 0.49$  vs  $2.28 \pm 0.45$ ) and there was no difference in sedation score between two groups at 60 minutes after surgery.

This was consistent with the results obtained by

- 1) Cr patel et al(2012)studied the effect of intravenous infusion of dexmedetomidine of on perioperative haemodynamic changes and post operative recovery and found that post operative sedation was significantly higher in dexmedetomidine group than control group.

### **TIME FOR RESCUE ANALGESIA**

In our study,the time for rescue analgesia was significantly longer in dexmedetomidine group ( $64.4 \pm 7.91$  minutes) than esmolol group( $40.64 \pm 5.56$  minutes).

This was consistent with the results obtained by

- 1) Gurbet et al(2006) made a study on intraoperative use of dexmedetomidine on post operative analgesic requirements and found that requirement of morphine in the post operative period was lower in dexmedetomidine group and concluded that dexmedetomidine provides excellent post operative analgesia without increasing side effects.
- 2) Gupta et al(2013) studied the effect on intraoperative dexmedetomidine on post operative recovery in children undergoing spinal surgery and found that post operative agitation

and pain scores were lower in dexmedetomidine group and requirement of analgesic's and time for rescue analgesia was longer in dexmedetomidine group.

### **INTRA OPERATIVE SERUM CORTISOL**

In our study, there was no significant differences between two group's with respect to intraoperative serum cortisol levels.

This is consistent with results obtained by

- 1) Sarpkaya et al (2010) studied the effects of perioperative use of dexmedetomidine on surgical stress response in hypertensive patients and found that serum cortisol levels were lower in dexmedetomidine group than control group.

## SUMMARY

This study was undertaken to compare dexmedetomidine with esmolol in controlled hypotension for functional endoscopic sinus surgery.

A total of 50 patients were randomly allocated into two groups .Group D received dexmedetomidine at a loading dose of 1 microgram/kg followed by infusion at a dose of 0.4-0.8 microgram/kg/hr.

Group E received esmolol at a loading dose of 1mg/kg followed by infusion at 0.4-0.8 mg /kg/hr.

Various parameters like dose of propofol, intraoperative heart rate and blood pressure, quality of surgical field, emergence time, post anaesthesia recovery score, sedation score, intra operative serum cortisol, time for rescue analgesia was recorded.

The observation noted were as follows

- The induction dose of propofol was lower in dexmedetomidine group than esmolol ,which was statistically significant.
- Heart rate and blood pressure was significantly lower in dexmedetomidine group than esmolol group at 5 and 10 min's after stoppage of infusion, at end of surgery and after extubation.
- The quality of surgical field was comparable between two groups.
- Emergence time was significantly prolonged in dexmedetomidine group than esmolol group.
- Time to achieve post anaesthesia recovery score (modified aldrete score>9)was significantly higher in dexmedetomidine group than esmolol group.
- Sedation score's were higher in dexmedetomidine group than esmolol group at 15 min's and 30 min's after extubation.
- Time for rescue analgesia was significantly higher in dexmedetomidine group than esmolol group.
- Intra operative serum cortisol level's were comparable between two group's.
- No adverse complications were noted in both group's.

## **CONCLUSION**

The observation of this study is that both esmolol and dexmedetomidine can be effectively used in induced hypotension for functional endoscopic sinus surgery. Both groups were comparable with respect to quality of surgical field and intraoperative haemodynamics.

This study also shows that dexmedetomidine has added advantage of sedative, analgesic and reduced requirement of anaesthetic agents when compared with esmolol. Post operative recovery scores and sedation scores were significantly higher and time required for rescue analgesia was also prolonged with dexmedetomidine than esmolol.



## BIBLIOGRAPHY

1. Miller's Anaesthesia -7<sup>th</sup> edition
2. Anaesthesiologist;s manual of surgical procedures by Richard A Jaffe,Stanley Samuels, Clifford Schmeising.
3. Boezaart AP, Van Der Merwe J , Coatzee A.-Comparison of sodium nitroprusside and esmolol induced controlled hypotension for functional endoscopic sinus surgery. Canadian Journal of Anaesthesia 1995 may,42(5pt1):373-6.
4. Koi IO, Kayqusuz K, Yildirim A, Dogan M, Gursoy S-Controlled hypotension with desflurane combined with esmolol (or) dexmedetomidine during tympanoplasty in adults, a double blinded randomized controlled trial.-Current therapeutic research.2009jan 70(3) 197-208.
5. Turan G,Dincer E-Comparison of dexmedetomidine,remifentanil and esmolol in controlled hypotensive anaesthesia-Internet journal of anaesthesia 2007 volume 17 number 2.
6. MA Basyoni, Abdul Mohsin J, Alsarraf- Induced hypotension for nasal surgery remifentanil versus esmolol- Journal of American academy of otolaryngology head and neck surgery 2013 volume 149 no 2.
7. YJ lim,C S Kim,J S Bahk,BM Ham-Clinical trial of esmolol induced controlled hypotension with (or) without acute normovolemic hemodilution in spinal surgery-Acta anaesthesiologica scandinavica 2003 jan, volume 47 pages 74-78.

8. Celebi N, Artukoglu F ,Savicaoglu F-Effect of hypotensive anaesthesia on Cognitive function ,a comparison of esmolol and remifentanil during tympanoplasty-Saudi medical journal sep 2007 ,28(9)-1357-66.
9. Aylu Guney,Fatma NurKaya,Gurnet-Comparison of esmolol to nitroglycerine in controlled hypotension during nasal surgery-Eurasian journal of medicine 2012,44:99-105.
10. Eun HH, Hwang HY, Ryu HH, Lee YW-Comparison of sodium nitroprusside and esmolol induced hypotension for total hip arthroplasty – Korean journal of anaesthesia 1997 aug 33(2):324-329.
11. Wylie Churchill-davidson's a practice of anaesthesia 7<sup>th</sup> edition-Control of circulation page 203-219.
12. Gokce BM,Karabiyi kL,Karadenizli Y-Hypotensive anaesthesia with esmolol assessment of hemodynamics,consumption of anaesthetic drugs and recovery-Saudi medical journal 2009 vol 30(6) 771-777.
13. Kumsuk park, Jong su kim-Effects of hydralazine pretreatment on esmolol induced controlled hypotension during spine surgery-Korean journal of anaesthesia- june 2006 vol 50 ,no 6.
14. Himanshu dongre,Vijay sharma,Aditi dongre-The efficacy of esmolol and nitroglycerine in creating dry operating field by producing controlled hypotension in spinal surgeries-IOSR journal of pharmacy july 2012,volume2,pp 26-33.
15. Zynel abiding erbesler, Nurten bakan –Comparison of effects of esmolol and dexmedetomidine on clinical course and cost for controlled hypotensive anaesthesia-Turkish journal of anaesthesia 2013 41:page 156-61.

16. Mengitsu AM, Wolf MW, Boldt J-Influence of controlled hypotension with esmolol and sodium nitroprusside on natriuretic peptides undergoing endonasal sinus surgery-European journal of anaesthesia June 2007 pp 529-534.
17. Akcan akkayan, Gmityasav tekelioglu-Comparison of effects of magnesium sulphate and dexmedetomidine on surgical vision quality on endoscopic sinus surgery.randomized clinical study-Brazilian journal of anaesthesiology 2014 jan
18. Shen PH , Weitzel EK , Lai JT-Intravenous esmolol infusion improves surgical Field's during sevoflurane anaesthetized endoscopic sinus surgery.a double blind , randomized placebo, controlled trial-American journal of rhinology and allergy nov/dec 2001 volume 25, number 6, pp 208-211.
19. Abdulla aydinozcan, Yaman ozyurt, Aakan erkul-Dexmedetomidine versus remifentanyl for controlled hypotensive anaesthesia in functional endoscopic sinus surgery-Turkish journal of anaesthesia 2012 :40(5):257-61.
20. Farah nasreen , Shahjahan bano, Rashid mansoor khan-Dexmedetomidine used to provide hypotensive anaesthesia during middle ear surgery-Indian journal of otolaryngeal head and neck surgery jul/sep 2009 61: 205-207.
21. Bayazit dikmen, Fazilet sahin, Bilse nornek -Dexmedetomidine for controlled hypotension in middle ear surgery with low flow anaesthesia-International journal of advanced otology 2010 6(3) 331-336.
22. Richa F, Yaziqi A, Yazbeck P-Comparison between dexmedetomidine and remifentanyl for controlled hypotension during tympanoplasty-European journal of anaesthesiology 2008 pg 369-74.

23. Malhotra SK ,Kumar S, Arora S, Gupta A-Effect of dexmedetomidine on quality of surgical field and requirement of hypotensive agents during endoscopic sinus surgery under general anaesthesia-European journal of anaesthesiology june 2013 volume 30 pg 27-37.
24. Guang jiecao,Ying yang,Bing wang-Faesibility of dexmedetomidine with sevoflurane for controlled hypotension in endoscopic sinus surgery- Medical journal of chinese people liberation army 2012 volume 37 ,no 1.
25. Sunil chiruvella, Balaji bonthu ,Venkatasiva,Dorababu S-Controlled Hypotensive anaesthesia with dexmedetomidine for FESS .a prospective randomized double blind study -Journal of evolution of medical and dental sciences august 2014 ,volume 3 page 9556-9563.
26. Gurbet A,Basavanmoqole ,Turker G-Intraoperative infusion of Dexmedetomidine reduces perioperative analgesic requirements –Canadian journal of anaesthesia2006 jul :33(7) 646-51.
27. Kim JM,Lee JH,Song BM,Koo BN-Efficacy of intraoperative dexmedetomidine on emergence agitation and quality of recovery after nasal surgery-British journal of anaesthesia 2013 aug :111(1) 222-8.
28. Guray demir,Kulay even-Effect of dexmedetomidine on propofol amount and hemodynamic parameters at anaesthesia induction –Medical journal of Bakirkoy,2009 ,volume 5,number 2.
29. Fazilet sahin,Devpi lderen,Gulayer dogan–Comparison of dexmedetomidine And alfentanil during middle ear surgery-International journal of advanced otology2011:7(2) pg 225-233.

30. CRPatel ,Engineer SR,Shah BJ ,Madhu S-Effect of intravenous infusion of dexmedetomidine on perioperative haemodynamic changes and post operative recovery-A study with entropy analysis 2012 ,volume 56,page 542-546.
31. Suvadeep , Chakraborty, Jayanta-Effect of dexmedetomidine infusion on Propofol requirement for maintenance of optimum depth of anaesthesia during elective spine surgery-Indian journal of anaesthesia 2013,volume 57 ,page 358-363.
32. Gupta N,Ram GP, Prabakar H-Effect of intraoperative dexmedetomidine on post op recovery profile of children undergoing surgery for spinal dysraphism-Journal of neurosurgical anaesthesiology 2013 jul 25(3):271-278.
33. Kana H,Kunisava J, Kurosava A-Effect of dexmedetomidine on anaesthetic requirements in cardiovascular surgery-Japanese journal of anaesthesiology 2009dec 58(11)1496-1500.
34. Sarpkaya,Karaaslan K, Kocoglu H, Colak C-Effects of perioperative use of dexmedetomidine on hemodynamic parameters and surgical stress response in chronic hypertensive patients –European journal of anaesthesiology 12 june2012 ,volume 27,pg 86-89.

# PROFORMA

Name:

Age:

Sex:

Weight:

Diagnosis:

Surgery:

MPC:

ASA PS class:

Group:

parameters		
Dose of propofol		
Heart rate	Pre op	
	After induction	
	15 min of surgery	
	30 min of surgery	
	45 min of surgery	
	5 min after stop	
	10 min after stop	
	At end of surgery	
	After extubation	
Blood pressure	Pre op	
	After induction	
	15 min of surgery	
	30 min of surgery	
	45 min of surgery	
	5 min after stop	
	10 min after stop	
	At end of surgery	
	After extubation	
Intra op plasma cortisol		
Quality of surgical field		
Emergence time		
Post op recovery score		
Sedation score	15 minutes	
	30 minutes	
	60 minutes	
Time for rescue analgesic		

MASTER CHART GROUP D

group	Name	age	sex	wt	Sbp preop	Dbp preop	Sbp after int	Dbp after int	sbp15mi	dbp15mi	sbp30mi	dbp30 min	sbp45mi	dbp45mi	sbp5 min af stop	dbp5min af stop	sbp10mi af stop	dbp10 min af stop	Sbp at end	Dbp at end	Sbp af ex	Dbp af ex
1	Perumal	38	1	50	120	72	108	70	90	58	90	56	90	58	94	60	96	62	98	70	102	71
1	Thamilarasi	31	2	65	120	70	110	71	93	52	90	54	90	58	98	64	96	60	100	62	103	63
1	Verrakannan	21	1	50	125	60	106	70	98	54	86	52	80	53	98	58	100	60	96	70	98	74
1	Gajendran	42	1	70	120	70	110	70	96	58	80	52	80	54	92	60	93	65	94	66	97	70
1	Rukumani	36	1	40	116	78	105	70	88	55	88	56	80	58	94	63	93	60	96	66	98	63
1	Raguram	26	1	60	120	70	106	70	89	58	85	52	84	58	90	66	94	68	96	65	99	66
1	Latha	30	2	50	110	78	100	74	90	50	88	52	86	50	90	54	93	62	98	60	103	70
1	Sasikala	34	2	55	106	72	102	66	91	58	80	58	88	54	90	62	96	67	92	64	105	70
1	Thenmozhi	40	2	45	120	70	120	68	99	52	90	56	80	54	96	62	95	60	93	60	103	63
1	Sathish	30	1	75	114	70	110	70	84	58	88	56	85	50	90	62	93	60	98	62	98	66
1	Karthick	28	1	65	120	71	117	66	82	56	84	56	88	54	94	64	98	60	102	62	103	63
1	Manimaran	45	1	55	120	72	111	60	89	58	86	52	90	56	92	58	90	57	94	69	98	66
1	Dinesh	23	1	60	117	80	110	70	93	58	90	60	88	54	90	62	93	63	98	60	101	65
1	Suganya	27	2	50	110	70	108	70	90	60	85	54	92	56	94	60	98	61	90	64	102	66
1	Vijayalakshmi	33	2	40	120	68	112	66	92	54	80	58	92	54	95	58	90	50	94	63	104	65
1	Subash	27	1	60	120	76	120	72	84	58	80	54	86	60	90	64	93	70	90	66	96	68
1	Nithya	30	2	50	120	78	116	70	83	60	84	60	88	63	92	66	97	62	94	60	97	64
1	Geetha	36	2	65	117	70	112	72	86	54	84	60	86	64	90	68	93	60	96	63	98	68
1	Vularmani	48	2	50	112	80	104	72	90	58	82	60	90	56	92	60	95	63	94	64	98	63
1	Umakannan	40	1	60	114	70	107	66	94	63	90	60	82	64	94	60	97	60	98	68	101	70
1	Lakshmi	37	2	50	126	80	112	66	84	60	85	63	88	60	90	62	92	60	96	64	98	67
1	Senthamarai	43	2	55	120	79	114	72	88	56	84	58	88	70	93	72	98	66	90	70	106	73
1	Kalyani	23	2	60	121	84	107	70	96	58	90	62	94	56	95	55	95	60	94	68	110	74
1	Karunakaren	35	1	70	109	70	104	76	90	58	90	62	94	58	91	55	90	56	94	60	108	74
1	Kannan	24	2	50	117	85	112	80	84	62	86	56	90	60	88	57	90	58	98	66	104	65



Pre op hr	Af intubation	Hr 15 min	Hr 30 min	Hr 45 min	Hr 5 min af stop	Hr 10 min af stop	At end of surgery	After extubation
86	72	62	62	56	56	62	64	67
84	70	62	62	56	56	64	66	66
86	66	60	66	54	50	63	68	68
84	66	60	68	54	54	64	65	66
80	70	62	70	56	60	68	70	71
70	70	60	60	56	58	63	65	64
76	66	62	60	58	58	61	64	66
80	66	62	62	56	54	62	65	67
84	70	66	66	58	60	67	69	70
90	70	62	70	56	60	71	73	71
86	66	60	74	58	60	75	77	78
82	66	60	63	58	60	60	62	63
76	66	60	58	55	55	60	64	65
86	70	60	67	58	65	66	68	69
68	60	56	52	50	50	54	56	60
74	66	60	61	59	60	61	62	65
70	60	56	54	51	53	55	57	60
80	72	64	64	58	64	65	66	68
78	66	60	66	56	69	70	73	71
88	66	60	69	58	72	71	72	74
92	66	60	71	58	68	70	73	75
90	66	60	67	60	71	72	74	76
76	66	60	58	56	56	57	59	63
94	70	62	71	60	72	73	74	78
92	72	64	70	58	70	74	75	78

Average category scale	Dose of propofol	Emergence time	Postanesthesia recovery score	Rescue analgesia	S 15 mins	S 30 mins	S 60 mins	Serum cortisol
2	70	12	15	60	3	3	2	4
2	80	12	15	65	3	2	2	3
2	60	12	14	55	3	3	2	3
2	80	13	14	65	3	3	2	4
2	50	14	16	70	3	3	2	4
2	75	15	17	75	3	3	2	5
2	65	13	14	80	3	3	2	5
2	65	16	16	65	3	3	2	4
2	60	11	15	60	3	2	2	4
2	90	13	14	60	2	3	2	5
2	80	13	15	70	3	3	2	4
2	75	14	16	75	2	2	2	3
2	70	14	16	62	3	3	2	4
2	60	13	11	45	3	3	2	4
2	60	13	18	80	2	2	2	4
2	70	14	15	55	3	3	2	5
2	60	14	16	60	3	2	2	5
2	70	15	16	65	3	3	2	4
2	70	14	17	70	2	2	2	4
2	65	12	11	60	3	3	2	4
2	60	13	16	65	2	2	2	4
2	70	13	18	65	3	3	2	4
2	70	13	17	60	2	2	2	4
2	80	14	18	63	3	3	2	3
2	70	13	18	60	3	2	2	4

GROUP E

group	Name	age	sex	wt	Sbp pre op	Dbp pre op	Sbp after intu	Dbp after intubation	Sbp 15 min	Dbp 15 min	Sbp 30 min	Dbp 30 min	Sbp 45 min	Dbp 45 min	Sbp 5 min af stop	Dbp 5 min af stop	Sbp 10 min af stop	Sbp at end	Dbp at end	Dbp at end	Sbp af ex
2	Rajasekar	21	1	85	120	80	110	70	96	52	90	50	94	54	106	57	107	70	108	114	78
2	Bala	50	2	40	120	70	100	74	92	58	88	54	91	54	109	62	106	64	112	120	74
2	sivakumar	21	1	50	130	80	110	76	96	56	94	60	93	62	110	63	114	71	116	126	80
2	Suganthi	27	2	60	126	74	116	76	100	50	93	58	90	54	104	70	116	81	114	124	80
2	Mythili	23	2	60	117	80	112	76	104	76	91	60	88	60	108	74	117	83	124	120	84
2	Tamilarasi	31	2	65	128	76	120	78	108	86	88	58	84	62	112	76	120	76	126	118	81
2	Jothi	31	2	54	108	76	106	80	98	58	99	60	93	62	120	72	114	76	117	118	80
2	Mohanapriya	22	2	45	114	72	110	72	90	56	80	58	84	62	107	78	108	82	110	100	70
2	Revathi	40	2	50	120	80	110	70	90	62	93	64	90	66	106	84	110	82	111	110	84
2	Suresh	27	1	60	116	73	114	74	96	63	90	60	92	62	102	78	106	76	121	124	82
2	Ramesh	30	1	55	127	90	120	76	86	60	85	65	84	60	109	70	114	76	110	116	83
2	Ramkumar	34	1	60	134	94	130	80	94	70	90	62	94	68	118	80	111	86	112	114	82
2	Ratnavel	47	1	68	130	82	126	76	90	72	92	67	90	61	112	72	114	76	111	120	86
2	Chitra	26	2	43	121	70	120	88	87	57	90	56	94	60	116	71	118	70	110	122	72
2	Shobana	28	2	48	114	80	110	70	88	53	92	58	96	60	121	70	118	78	120	124	84
2	Sukumar	22	1	50	112	68	110	62	92	51	90	56	88	60	108	81	126	70	124	126	79
2	Ravi	33	1	55	126	70	112	70	94	58	93	62	95	66	113	70	120	76	126	120	76
2	Ganesh	38	1	60	120	77	114	72	91	60	96	62	94	63	111	74	114	80	120	124	80
2	Selvakumar	40	1	53	114	84	106	80	95	56	90	62	93	60	110	71	118	81	123	116	80
2	Prabu	26	1	70	120	86	120	70	96	66	93	62	90	64	126	76	120	81	114	115	84
2	Karpagam	38	2	62	128	78	118	71	85	60	90	64	92	60	110	71	126	84	125	106	84
2	Praveen	21	1	68	132	80	120	76	87	64	88	62	85	60	114	76	116	80	110	126	83
2	Senthil	35	1	60	130	84	114	70	93	68	83	68	90	60	112	62	114	76	116	120	72
2	Sridhar	38	1	50	136	88	120	70	90	62	90	62	94	63	126	78	124	84	120	120	84
2	Sakthivel	47	1	55	120	70	110	62	89	63	89	63	95	66	114	82	112	76	110	113	80

Preop hr	Af intubation	Hr 15 min	Hr 30 min	Hr 45 min	Hr 5 min af stop	Hr 10 min af stop	At end of surgery	After extubation
92	72	64	70	58	70	74	75	78
84	68	64	60	60	68	72	72	76
90	70	62	60	58	66	70	74	74
96	80	72	70	72	80	86	86	88
90	76	70	68	66	70	74	76	80
80	70	64	62	62	68	70	72	80
86	76	66	64	66	70	74	74	90
92	80	68	62	60	64	68	70	82
72	60	52	52	50	58	64	66	76
86	70	66	62	60	64	66	70	78
78	62	56	52	50	56	62	64	70
98	72	60	58	56	62	66	68	76
104	86	72	68	64	70	76	74	80
78	64	56	54	52	60	66	64	78
84	70	60	56	52	56	60	64	70
90	72	62	60	58	66	70	72	76
96	80	72	66	64	70	76	78	86
100	84	76	70	70	76	82	84	90
92	72	68	64	64	72	78	80	90
80	64	60	54	52	60	64	68	80
86	66	58	52	54	62	66	70	82
84	60	56	54	56	62	68	72	82
78	64	60	56	54	64	70	70	80
90	75	70	66	68	76	80	84	90
96	80	74	70	66	74	80	86	90
78	62	56	54	52	58	66	70	84

Average category scale	Dose of propofol	Emergence time	Postanesthesia recovery score	Rescue analgesia	S 15 mins	S 30 mins	S 60 mins	Serum cortisol
2	70	13	18	60	3	2	2	4
2	150	5	10	35	2	2	2	3
2	70	7	11	40	3	3	2	4
2	75	7	11	42	2	2	2	5
2	100	8	12	30	2	2	2	3
2	90	6	9	45	3	3	2	4
2	100	9	11	40	2	2	2	4
2	90	8	10	50	2	2	2	3
2	80	6	11	35	2	2	2	3
2	80	7	11	45	3	3	2	3
2	110	8	12	40	2	2	2	3
2	80	9	12	46	2	2	2	4
2	90	6	13	48	2	2	2	4
2	100	6	11	35	3	3	2	4
2	70	8	10	30	3	2	2	3
2	90	9	11	40	2	2	2	3
2	90	10	11	45	2	2	2	3
2	100	8	12	40	2	2	2	3
2	90	7	12	35	3	3	2	4
2	80	8	13	45	2	2	2	4
2	120	8	12	50	2	2	2	4
2	100	9	11	40	3	2	2	4
2	120	10	11	45	2	2	2	4
2	100	9	10	40	3	3	2	4
2	90	7	9	40	2	2	2	3
2	90	8	10	35	3	3	2	5